

Dickerson, B.C., Salat, D.H., Greve, D.N., Chua, E.F., Rand-Giovannetti, E., Rentz, D.M., Bertram, L., Mullin, K., Tanzi, R.E., Blacker, D., & Albert, M.S. (2005) *Increased hippocampal activation in mild cognitive impairment compared to normal aging and AD. Neurology, 65(3): 404-411.*

Yassa, M.A., Stark, S.M., Bakker, A., Albert, M.S., Gallagher, M., Stark, & C.E. (2010) *High-resolution structural and functional MRI of hippocampal CA3 and dentate gyrus in patients with amnesic Mild Cognitive Impairment. Neuroimage 51: 1242-1252.*

Putchá, D., Brickhouse, M., O'Keefe, K., Sullivan, C., Rentz, D., Marshall, G., Dickerson, B., & Sperling, R. (2011) *Hippocampal hyperactivation associated with cortical thinning in Alzheimer's disease signature regions in non-demented elderly adults. The Journal of Neuroscience 31: 17680-17688.*

Huijbers, W., Mormino, E.C., Schultz, A.P., Wigman, S., Ward, A.M., Larvie, M., Amariglio, R.E., Marshall, G.A., Rentz, D.M., Johnson, K.A., & Sperling, R.A. (2015) *Amyloid- β deposition in mild cognitive impairment is associated with increased hippocampal activity, atrophy and clinical progression. Brain 138: 1023-1035.*

Leal, S.L., Landau, S.M., Bell, R.K., & Jagust, W.J. (2017) *Hippocampal activation is associated with longitudinal amyloid accumulation and cognitive decline. Elife 6: e22978.*

Marks, S.M., Lockhart, S.N., Baker, S.L., & Jagust, W.J. (2017) *Tau and β -amyloid are associated with medial temporal lobe structure, function and memory encoding in normal aging. The Journal of Neuroscience: 3769-16.*

Huijbers, W., Schultz, A.P., Papp, K.V., LaPoint, M.R., Hanseeuw, B., Chhatwal, J., Hedden, T., Johnson, K.A. & Sperling, R. (2019) *Tau accumulation in clinically normal older adults is associated with hippocampal hyperactivity. The Journal of Neuroscience. 39: 548-556.*

Berron, D., Cardenas-Blanco, A., Bittner, D., Metzger, C.D., Spottke, A., Heneka, M.T., Liessbach, K., Schneider, A., Teipel, S.J., Wagner, M., Speck, O., Janssen, F., & Duzel, E. (2019) *Higher CSF Tau levels are related to hippocampal hyperactivity and object mnemonic discrimination in older adults. (2019) The Journal of Neuroscience. 39: 8788-8797.*

Richter, N., Bischof, G., Dronse, J., Nellessen, N., Neumaier, B., Langen, K.-J., Drzezga, A., Fink, G.R., van Eimeren, T., Kukolja, J., & Onur, O.A. (2020) *Entorhinal tau predicts hippocampal activation and memory deficits in Alzheimer's disease. Journal of Alzheimer's Disease. 78: 1601-1604.*

Palmqvist S, Janelidze S, Quiroz YT, Zetterberg H, Lopera F, Stomrud E, Su Y, Chen Y, Serrano GE, Leuzy A, Mattsson-Carlgrén N, Strandberg O, Smith R, Villegas A, Sepulveda-Falla D, Chai X, Proctor NK, Beach TG, Blennow K, Dage JL, Reiman EM, Hansson O. (2020) *Discriminative Accuracy of Plasma Phospho-tau217 for Alzheimer Disease vs Other Neurodegenerative Disorders. JAMA 324(8):772-781.*

Mattson-Carlgrén, N., Janelidze, S., Palmqvist, S., Cullen, N., Svenningsson, A.L., Strandberg, O., Mengel, D., Walsh, D.M., Stomrud, E., Dage, J.L., Hansson, O. (2020). *Longitudinal plasma p-tau217 is increased in early stages of Alzheimer's disease. Brain. 143: 3234-3241.*
